

MERIT BADGE SERIES



ASTRONOMY



BOY SCOUTS OF AMERICA®



STEM-Based

BOY SCOUTS OF AMERICA
MERIT BADGE SERIES

ASTRONOMY



"Enhancing our youths' competitive edge through merit badges"



BOY SCOUTS OF AMERICA®

Requirements

Always check www.scouting.org for the latest requirements.

1. Do the following:
 - a. Explain to your counselor the most likely hazards you may encounter while participating in astronomy activities, and what you should do to anticipate, help prevent, mitigate, and respond to these hazards.
 - b. Explain first aid for injuries or illnesses such as heat and cold reactions, dehydration, bites and stings, and damage to your eyes that could occur during observation.
 - c. Describe the proper clothing and other precautions for safely making observations at night and in cold weather.
 - d. Explain how to safely observe the Sun, objects near the Sun and solar eclipses.
2. Explain what light pollution is and how it and air pollution affect astronomy.
3. With the aid of diagrams (or real telescopes if available), do each of the following:
 - a. Explain why binoculars and telescopes are important astronomical tools. Demonstrate or explain how these tools are used.
 - b. Describe the similarities and differences of several types of astronomical telescopes, including at least one that observes light beyond the visible part of the spectrum (i.e., radio, X-ray, ultraviolet, or infrared).
 - c. Explain the purposes of at least three instruments used with astronomical telescopes.
 - d. Describe the proper care and storage of telescopes and binoculars both at home and in the field.

For requirement
4, if instruction
is done in a
planetarium,
Scouts must still
identify the
required stars and
constellations
under the natural
night sky.

4. Do the following:
 - a. Identify in the sky at least 10 constellations, at least four of which are in the zodiac.
 - b. Identify in the sky at least eight conspicuous stars, five of which are of magnitude 1 or brighter.
 - c. Make two sketches of the Big Dipper. In one sketch, show the Big Dipper's orientation in the early evening sky. In another sketch, show its position several hours later. In both sketches, show the North Star and the horizon. Record the date and time each sketch was made.
 - d. Explain what we see when we look at the Milky Way.
5. Do the following:
 - a. List the names of the five most visible planets. Explain which ones can appear in phases similar to lunar phases and which ones cannot, and explain why.
 - b. Using the internet (with your parent's permission) and other resources, find out when each of the five most visible planets that you identified in requirement 5a will be observable in the evening sky during the next 12 months. Then compile this information in the form of a chart or table.
 - c. Describe the motion of the planets across the sky.
 - d. Observe a planet and describe what you saw.
6. Do the following:
 - a. Sketch the face of the Moon and indicate at least five seas and five craters. Label these landmarks.
 - b. Sketch the phase and position of the Moon, at the same hour and place, for four nights within a one week period. Include landmarks on the horizon such as hills, trees, and buildings. Explain the changes you observe.
 - c. List the factors that keep the Moon in orbit around Earth.
 - d. With the aid of diagrams, explain the relative positions of the Sun, Earth, and the Moon at the times of lunar and solar eclipses, and at the times of new, first-quarter, full, and last-quarter phases of the Moon.

7. Do the following:
- Describe the composition of the Sun, its relationship to other stars, and some effects of its radiation on Earth's weather and communications.
 - Define sunspots and describe some of the effects they may have on solar radiation.
 - Identify at least one red star, one blue star, and one yellow star (other than the Sun). Explain the meaning of these colors.
8. With your counselor's approval and guidance, do ONE of the following:
- Visit a planetarium or astronomical observatory. Submit a written report, a scrapbook, or a video presentation afterward to your counselor that includes the following information:
 - Activities occurring there
 - Exhibits and displays you saw
 - Telescopes and other instruments being used
 - Celestial objects you observed
 - Plan and participate in a three-hour observation session that includes using binoculars or a telescope. List the celestial objects you want to observe, and find each on a star chart or in a guidebook. Prepare a log or notebook. Discuss with your counselor what you hope to observe prior to your observation session. Review your log or notebook with your counselor afterward.*
 - Plan and host a star party for your Scout troop or other group such as your class at school. Use binoculars or a telescope to show and explain celestial objects to the group.
 - Help an astronomy club in your community hold a star party that is open to the public.
 - Personally take a series of photographs or digital images of the movement of the Moon, a planet, an asteroid, meteor, or a comet. In your visual display, label each image and include the date and time it was taken. Show all positions on a star chart or map. Show your display at school or at a troop meeting. Explain the changes you observed.
9. Find out about three career opportunities in astronomy. Pick one and find out the education, training, and experience required for this profession. Discuss this with your counselor, and explain why this profession might interest you.

*To complete requirement 8b, you may use the Scout Planning Worksheet: troopleader.org/wp-content/uploads/2016/03/512-505_16_Wksht_WEB.pdf

Contents

Introduction: Reach for the Stars 7

Getting Ready to Observe. 8

Stars and Constellations. 13

The Planets and Our Solar System 27

The Moon—Our Nearest Neighbor. 39

The Sun and Stars 47

Tools of the Trade 55

Light Pollution. 67

You and Astronomy 69

Careers in Astronomy 73

Astronomy Resources 77





Introduction: Reach for the Stars

Spectacular shows can be seen in the night sky. From Earth, we can see *nebulae*, or giant clouds of gas and dust. We can see stars dying and exploding. Sometimes it is rare to see sights like meteor showers or dramatic eclipses. But on any clear night of the year, you usually can see the Moon and, if your skies are clear and mostly free from light pollution, you can see our whole galaxy stretching across the sky.


In learning about astronomy, you will study how activities in space affect your own planet. Your daily schedule—daytime and nighttime—is dictated by Earth’s rotation. Earth’s orbit around the Sun takes a year and gives us the changing seasons. The Sun’s energy affects the weather, and the gravitational pull between Earth and the Moon creates and controls ocean tides.

We still have much to learn about space. It wasn’t until 2011 that NASA’s Kepler spacecraft discovered two Earth-sized planets beyond the solar system. Instruments such as the Hubble Space Telescope enable us to see visions today that earlier observers could not have imagined.


We know much about space and its effects on us, but there is a great deal we don’t yet know. Someday you might be among the scientists or the serious amateur observers who make important new discoveries.

An *axis* is the imaginary line around which a rotating body, such as Earth, turns. Earth’s axis is a straight line between the geographic (not magnetic) north and south poles. Earth’s *orbit* is the curved path it follows on its yearly trip around the Sun.

Just as you enjoy looking up at the stars and finding the Big Dipper, ancient stargazers had fun naming the patterns of stars they saw every night. The word *astronomy* literally translates to “naming the heavens.”



REVOLUTION



ROTATION



Getting Ready to Observe

Know how to protect yourself before you go out in search of the next great astronomical discovery.

Protect Your Body

When you go stargazing, dress appropriately for the weather—hot or cold. To help protect yourself against bites and stings when you go outside, wear clothes that cover all exposed skin, and be sure to button and tuck them in. Wear shoes or boots—not sandals—and socks. Insect repellent sprayed on your body, clothing, and shoes provides added protection.

Many insect repellents contain chemicals that may corrode plastic. Whenever you are applying insect repellent, do so where it won't affect your equipment. Afterward, clean your hands before handling a telescope.



It takes about 30 minutes for the human eye to adjust to darkness. To help preserve your night vision, use a red filter with your flashlight. Simply secure a piece of red cellophane over an ordinary flashlight.

Moving in the Dark

During the day before a stargazing outing, survey the area where you will be observing. Look for drop-offs, holes, crevices, or other objects like large rocks or tree roots that you might not see at night. Take care to avoid the hazards when you return at night to observe. Bring red-filtered flashlights to illuminate the area.


Protect Your Eyes

Never stare directly at or near the Sun, even for a few moments. **Never** look at the Sun through binoculars or a telescope unless your equipment has the proper solar filters. Looking directly at the Sun—even while wearing sunglasses—can cause permanent blindness or other damage to your eyes that might not be immediately noticeable. *The safest way to observe the Sun is indirectly by projection*, which is explained in the chapter called “The Moon—Our Nearest Neighbor.” Projection is the recommended method for safely viewing the sun. In any case, always do so only under the supervision of a knowledgeable adult.

Observing Objects Near the Sun

Mercury, the planet nearest the Sun, always lies close to the Sun in the sky. Look for Mercury only when the Sun’s disk is entirely below the horizon. Don’t risk getting the Sun in the view. This could cause permanent eye damage or blindness.

If you are observing the Sun and you or a companion experience eye discomfort, cover the eyes and seek medical attention immediately.



Heat Exhaustion and Heatstroke

Heat exhaustion occurs when the body overheats because its cooling methods fail. Watch for these signs: body temperature between 98.6 and 102 degrees; skin pale, clammy, and sweaty; nausea, dizziness, and fainting; pronounced weakness and tiredness; headache; muscle cramps. To treat heat exhaustion, have the victim lie down in a cool spot with the feet raised. Loosen the clothing. Apply cool, damp cloths to the skin or use a fan. Have the victim sip water.



Heatstroke (sunstroke) is life-threatening because the body's heat control system has been overworked and overwhelmed, resulting in its failure. Signs include body temperature above 102 degrees; red, hot, and dry skin; extremely rapid pulse; confusion or disorientation; fainting or unconsciousness; convulsions.

Cool the victim immediately. Place the victim in a cool spot face-up with head and shoulders raised. Remove outer clothing, sponge the bare skin with cold water, and soak underclothing with cool water. Apply cold packs, use a fan, or place the victim in a tub of

cold water. Dry the skin after the body temperature drops to 101 degrees. Obtain medical help immediately.

Sunburn

Sun exposure can catch you by surprise when you are outside, preoccupied with setting up your equipment or viewing a solar eclipse. Wear loose-fitting clothing that completely covers the arms and legs, and a brimmed hat. Apply sunscreen with a sun protection factor (SPF) of at least 15 to exposed skin. Don't forget your ears and the back of your neck. Reapply sunscreen often and as needed.

Cold-Related Problems—Hypothermia

Hypothermia occurs when the body's core temperature drops so low that it is no longer able to keep warm. The key to preventing hypothermia is to keep warm and stay dry, and—if you will be outside for extended periods—eat plenty of energy foods (nuts, dried fruits, peanut butter). Don't push yourself to a dangerous point of fatigue.

A person in the early stages of hypothermia may be shivering. As the victim gets colder, the shivering will stop. Other symptoms may include irritability, sleepiness, incoherence, disorientation, and the inability to reason or think clearly.

Immediately prevent further heat loss. Move the victim to a shelter, remove damp clothing, and warm the person with blankets until body temperature returns to normal. Cover the head with a warm hat or other covering, and offer hot drinks.

If the condition progresses, actively warm the victim's body. Place the victim into a sleeping bag with one or two other people. All should be stripped of most clothing so that skin-to-skin contact can hasten the warming—and perhaps save a life. Severe hypothermia requires immediate medical attention.

In cold weather, wear layers to keep in the heat.

Bring along a woolen cap, scarf, gloves, extra socks, and a coat. Keep your arms and legs covered.

Always carry rain gear.



Dehydration

Dehydration is caused by lack of water in the body.

A person who loses more water than consumed can become dehydrated—in hot *or* cold weather. Astronomical observers should stay well-hydrated while in the outdoors. Do not wait to drink until you feel thirsty.

Whenever you participate in astronomy activities, taking a few precautions can help make your outing a fun, safe, and comfortable one. By anticipating what can happen (such as exposure to cold or windy conditions), you can help prevent and mitigate (lessen the impact of) just about any type of mishap. You will also be in a better position to respond quickly and appropriately should a mishap occur. Remember to:

- Dress appropriately for the weather.
- Take and use the proper equipment, such as solar filters.
- Drink plenty of water; stay hydrated in warm *and* cool weather.



Circumpolar star chart

Stars and Constellations

People have always enjoyed studying the stars. To help them keep track of the stars, ancient Greeks creatively drew imaginary lines between stars to form images of mythological characters or other familiar creatures. These star groups are known as *constellations*. Today, astronomers recognize 88 constellations.

Star charts, which you can see on the following pages, help us find constellations. When using star charts, remember:

- a. The direction you are facing should be at the bottom of the chart. If you are facing south, hold the star chart with south at the bottom. If you are facing east, turn the chart so that east is at the bottom.
- b. On a star chart, east and west are reversed! This is because we are looking up into the sky rather than down onto Earth.
- c. Brighter stars have larger symbols on the map.
- d. The sky changes during the night, as the stars rise and set due to the rotation of Earth, and the sky changes with the seasons as Earth orbits the Sun. Thus, a star map is usually only accurate for a certain day and time.

A *star* is a hot, glowing ball of gas that produces energy by nuclear reactions in its core. In fact, our sun is a star.

A *constellation* is a pattern of stars in the sky.

Once you can identify a few of the bright constellations, such as Ursa Major or Orion, you can use them to find fainter and less obvious constellations.

Year-Round Constellations

Some constellations are visible any time of the year. These constellations are known as *circumpolar* because they never set below the horizon. As Earth rotates on its axis, the stars seem to circle a point called the celestial north pole. A moderately bright star, Polaris, currently lies close to the pole. Knowing the circumpolar constellations is a big help in locating Polaris in the sky.

Ursa Major. The Big Dipper is part of Ursa Major (the Great Bear), a constellation that includes a few less-visible stars. The outer stars on the bowl of the Big Dipper always point to Polaris. Every six hours, the Big Dipper appears to have moved a quarter of the way around Polaris.

Ursa Minor. The Little Dipper, with stars fainter than the Big Dipper, is part of the constellation Ursa Minor, or the Little Bear. Polaris is the end of the Little Dipper's handle.

Cassiopeia. If you follow the line from the middle star of the Big Dipper's handle through Polaris, you will find Cassiopeia. Its five main stars form an M when they're above the North Star and a W when they are below it.

Draco. Draco, the dragon, winds around Polaris between the Big and Little Dippers. None of its more than 80 visible stars is very bright; the four stars that form the dragon's head are easiest to see.

Seasonal Constellations

Some constellations are visible in the night sky only at certain times of the year. Twelve seasonal constellations known as zodiacal constellations are centered on the *ecliptic*, the path that the Sun, the Moon, and the planets all appear to follow through the sky. The zodiacal constellations you can see at night—Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpius, Sagittarius, Capricornus, Aquarius, and Pisces—change as Earth follows its yearly path around the Sun.

Spring Constellations

Leo. You can see Leo, the Lion, in the evening sky around 10:00 p.m., from January through June. You can use the Big Dipper to find Leo. Locate the two stars at the back of the bowl, next to the handle of the dipper. A line drawn from the star at the top of the bowl to the one at the bottom will point to Leo's brightest star, Regulus.

Some people believe the positions of the Sun, Moon, and planets among the zodiacal constellations affect human behavior. This belief is known as astrology. However, there is no scientific evidence to support this belief.

A *galaxy* is an enormous group of millions or billions of stars, bound together by gravity. Earth is in the Milky Way galaxy.

Virgo. Virgo, the Virgin, is the largest constellation and is visible from April through July. To use the Big Dipper to find Virgo, follow the arc of the handle to Arcturus, a bright star in the constellation Boötes (arc to Arcturus), then “drive a straight spike” down to the bright star Spica (spike to Spica), the bottom of Virgo’s Y shape.

Libra. Libra, the Balance Scale, is a late-spring and summer constellation with four fairly dim stars. It is one of the most difficult zodiacal star patterns to spot. Look halfway between Spica and Antares (in Scorpius).

Cancer. Cancer, the Crab, is visible from January to May, but it is the faintest constellation in the zodiac. To find it, follow a line from Leo’s Regulus to the Gemini constellation (a winter constellation described below). Cancer is between Leo and Gemini.

Summer Constellations

Lyra. You can see Lyra, the Lyre, from May through November. In late summer at about 10 p.m., if you look straight up, you may see a very bright blue-white star. This star, Vega, is Lyra’s brightest star.


Cygnus. Often called the Northern Cross, Cygnus the Swan is visible from June through November. Follow a line from Vega slightly east to another star, Deneb, which is Cygnus’ brightest star.

Of the conspicuous (easy-to-see) stars, only Sirius and Arcturus are brighter than Vega.

Aquila. South of Lyra and Cygnus is Aquila, the Eagle. Its brightest star is Altair, which forms the Summer Triangle with Deneb and Vega.

Part of Aquila lies along the path of the Milky Way where there are dense star fields.

Scorpius. In July and August, you can see Scorpius, the Scorpion, very close to the southern horizon. Its brightest star is Antares, a red supergiant star near the scorpion’s heart.



Summer star chart—around midnight in mid-July and 10 p.m. in mid-August

To locate the
Circlet, an
interesting star
group in Pisces,
look just south of
the Great Square
of Pegasus.

Sagittarius. In July and August you can see Sagittarius, the Archer, just east of Scorpius. Its main stars form a pattern that resembles a teapot. When you look at Sagittarius, you're looking toward the center of the Milky Way galaxy. Scorpius and Sagittarius may be difficult to view if you live in the North.

Autumn Constellations

Pegasus. Pegasus, the Flying Horse, is best seen from August through October, southeast of Cygnus. Part of the constellation is known as the Great Square, an easy shape to recognize.

Andromeda. Pegasus shares one of its stars with Andromeda, which is visible from September to January. Look eastward from the Great Square of Pegasus. The Andromeda galaxy—the most distant object visible to the naked eye—is visible on clear nights as a faint, misty spot in the Andromeda constellation.

Perseus. You can see Perseus in the autumn and winter, east of Andromeda. It lies between the constellations of Auriga (described later in this chapter) and Cassiopeia. Perseus contains the double star Algol. As the stars of Algol pass in front of each other, Algol appears alternately fainter and brighter.


A double star is a pair of stars that are close together and hold each other captive by the force of gravity. Each one orbits around the other. In most cases they are so close together, they look like single stars.

Aries. Three fairly bright stars make up the main part of Aries, the Ram. It appears south of Andromeda from October through March.

Pisces. The main part of Pisces, the Fishes, is a string of stars below Andromeda and Pegasus. Pisces appears in the sky from October to December, but it is faint and can be hard to find.

Capricornus. Look straight down from the star Altair in Aquila. Capricornus, the Sea Goat, is a faint constellation, but when visibility is good you can see it from August through October.

Aquarius. On dark, clear nights from August through October, you may be able to see Aquarius, the Water Bearer, south of Pegasus. One end of Aquarius stretches above Capricornus and the other is below the Circlet of Pisces.



Autumn star chart—around midnight in mid-October and 10 p.m. in mid-November

Betelgeuse, Orion's shoulder star, is so huge that if it were at the center of our solar system, its edges would extend beyond the orbit of Mars.

Winter Constellations

Orion. Orion, the Hunter, usually is easy to find from October through April. It is large and distinctive with two very bright stars—the reddish-orange giant Betelgeuse and bluish-white Rigel—as well as five other bright stars and several less visible ones. The Orion Nebula, a cloud of dust and gas several light-years across, is visible to the naked eye as a fuzzy patch just below Orion's belt.


Canis Major. Southeast of Orion is Canis Major, the Great Dog, which you can see from December through April. Sirius, the night sky's brightest star, is a part of Canis Major. Orion's three belt stars point downward toward Sirius.

Gemini. From December through May you can see Gemini, the Twins, which are above and to the left of Orion in the northern hemisphere. Two bright stars, Castor and Pollux, are the heads of the Gemini twins.

Auriga. Visible from November through April, Auriga, the Charioteer, is north of Orion. Capella, a double star in Auriga, is one of the brightest lights in the sky.

Taurus. Orion's belt stars point up toward Aldebaran, a bright orange-red star. Aldebaran is the eye of Taurus the Bull, a constellation you can see from November through March.

Taurus has two star groups: one a cluster of stars known as the Pleiades, and the other a larger, more scattered, V-shaped group known as the Hyades.



Winter star chart—around midnight in mid-January and 10 p.m. in mid-February

Star Light, Star Bright

Another word astronomers use to describe a star’s brightness is *magnitude*.

- *Absolute magnitude* refers to a star’s true brightness—that is, the brightness if all stars were viewed from the same distance.
- *Apparent magnitude* means the brightness of a star as we see it from Earth. Astronomers give a star a number to refer to this magnitude. The lower the magnitude number, the brighter the star. Stars that are brighter than a magnitude 1 have a zero or negative number. The unaided human eye can see stars with a magnitude number as high as 6.

Listed here are some of the brightest stars in the sky, the constellations where they are found, and their apparent magnitude.

Star	Constellation	Magnitude
Sirius	Canis Major	-1.46
Arcturus	Boötes	-0.04
Vega	Lyra	0.03
Capella	Auriga	0.08
Rigel	Orion	0.12
Procyon	Canis Minor	0.38
Betelgeuse	Orion	0.5
Altair	Aquila	0.77
Aldebaran	Taurus	0.85
Antares	Scorpius	0.96
Spica	Virgo	0.98
Pollux	Gemini	1.14
Deneb	Cygnus	1.25



If you live outside a major city, you probably have seen the Milky Way, which can be viewed without the aid of a telescope or binoculars.

The Milky Way

When we look at the Milky Way in the night sky, we see our own galaxy. Apart from other galaxies, everything we see in the night sky—the stars, star clusters, nebulae, the Sun, the Moon, and the planets—is part of the Milky Way. Just as its name implies, it appears as a whitish band of light. For observers in the northern hemisphere, it is easiest to see during the summer months in the constellations Aquila and Cygnus.

The Milky Way is our home galaxy, a gigantic disk of hundreds of billions of stars and vast quantities of dust and gas. It is about 100,000 light-years across and some 10,000 light-years thick at its center. Stars, dust, and gases fan out from the hub of this huge whirling disk in long, curving arms that make a spiral or pinwheel shape. Our solar system is about halfway out from the center, on a small spur between two of the arms.

Every one of the billions of galaxies in the universe has millions or billions of stars. The Milky Way appears to be an average spiral galaxy.



You can see another galaxy with the unaided eye on a clear night in the fall—the Andromeda Galaxy. It also is the most distant object you can see with the naked eye. It appears as a misty patch of light in the constellation of Andromeda. Astronomers call the Andromeda Galaxy “M31” and place it at about 2.5 million light-years from Earth. M31 appears to be half again as big as our galaxy—150,000 light-years across.


The Andromeda Galaxy is approximately 2.5 million light-years away. A *light-year* is the distance light travels in one year, equal to approximately 5.9 trillion miles.

Some galaxies can be seen through backyard telescopes at distances of 100 million light-years or more—distances almost unimaginable.

Using a Planisphere

A *planisphere* is a flat, circular map of the stars with an overlay that you can set to show what stars and constellations are visible at any time of the night and any night of the year. It can be adjusted to match the exact month, day, and hour when you are looking at the sky.





The Planets and Our Solar System

The planets of our solar system travel around the Sun, each in its own orbit. The four planets closest to the Sun—Mercury, Venus, Earth, and Mars—are called the terrestrial (earthlike) planets because they have solid, rocky surfaces. The four large planets beyond the orbit of Mars—Jupiter, Saturn, Uranus, and Neptune—are the gas giants, made mostly of hydrogen and helium, probably with no solid surfaces.

Mercury is the small, rocky planet nearest the Sun. It completes a trip around the Sun every 88 Earth-days, speeding in its orbit faster than any other planet in the solar system. If you could stand on Mercury, you would see a bloated sun creep at a snail's pace through a black sky. Mercury rotates extremely slowly on its axis—once in about 59 Earth-days. Because Mercury is so close to the Sun, temperatures on the planet's surface can reach a searing 800 degrees. Mercury's cratered surface looks much like the surface of Earth's moon.

Venus, second planet from the Sun, is only slightly smaller than Earth. The thick clouds of carbon dioxide and sulfuric acid that cover Venus reflect sunlight, making it bright but also trapping heat at the planet's surface. At 880 degrees F., it's hotter on Venus than on Mercury, even though Mercury is nearer the Sun. The thick, heavy atmosphere creates enormous pressure on Venus. Space probes that have landed there have lasted only a few hours before the pressure crushed them. Venus spins in the opposite direction of its orbit around the Sun.



Mercury



Venus is a scorched world with temperatures hot enough to melt lead.

**Earth****Mars****Jupiter**

Life on **Earth** survives beneath a thin layer of atmosphere that shelters us from the dangers of space. “Ocean” might be a more appropriate name for this planet, because oceans cover nearly 70 percent of Earth’s surface.

Mars, the Red Planet, is the fourth terrestrial planet. A day on Mars lasts 24 hours, 37 minutes—only slightly longer than an Earth-day. But Mars takes almost twice as long as Earth to orbit the Sun, going around once in 687 Earth-days.

Though Mars is a world with no rivers, it has winding valleys that look like dry riverbeds. Scientists believe that powerful floods once deluged Mars. Today the planet is too cold and its atmosphere too thin for liquid water to exist at the surface. In 2008, the Mars Phoenix lander confirmed the presence of ice on the surface, with more water frozen in the polar ice caps.

Jupiter, more than 300 million miles from Mars, is the largest planet in the solar system. If Jupiter were a hollow ball, more than a thousand Earths would fit inside. Despite its size, Jupiter spins faster than any other planet, rotating every 10 hours. However, it takes Jupiter nearly 12 Earth-years to orbit the Sun.

Like the other gas giants, Jupiter is mostly hydrogen and helium. The temperature at the top of its colorful clouds is minus 220 degrees. Down in the clouds, the temperature reaches 70 degrees at a depth where the atmospheric pressure is about 10 times as great as it is on Earth. Jupiter has at least 60 moons.

Three Earths could fit across Jupiter's Great Red Spot, a storm like a hurricane that has raged for centuries in Jupiter's southern hemisphere.



Saturn with its rings would just fit in the distance between Earth and the Moon.

Saturn

Saturn, the second largest planet, is the farthest from Earth of the five planets known to early stargazers. Galileo, in 1610, was the first astronomer to see Saturn and its rings through a telescope. Up to 1977, astronomers thought Saturn was the only planet with rings. We now know the other gas giants—Jupiter, Uranus, and Neptune—also have ring systems, but Saturn's is by far the largest, most visible, and most complex. Its rings are made mostly of ice. Saturn also has at least 80 moons. The largest, Titan, is a little bigger than Mercury and is blanketed with a thick, nitrogen-rich atmosphere.


Uranus (pronounced YOOR-un-nus), a large gas planet, blue-green in color, was discovered in 1781 by astronomer



Uranus

William Herschel; its rings were not discovered until 1977. Uranus is twice as far from the Sun as Saturn—so far away that it takes sunlight about 2 hours, 40 minutes to reach it. Uranus orbits the Sun tipped on its side, the result of what many astronomers think was a collision with a planet-sized body long ago.

On July 12, 2011, Neptune completed its first full orbit of the Sun since its discovery in 1846.



Neptune, the eighth planet from the Sun, is so far away that it takes 165 years to orbit the Sun. Like Earth, Neptune has seasons, but each season lasts for 41 Earth-years. Neptune is too distant to be seen with the unaided eye. In 1989, the Voyager 2 spacecraft tracked a large dark storm like a

hurricane in Neptune's southern hemisphere and discovered great geysers of nitrogen on Triton, Neptune's largest moon. Triton is one of the coldest bodies yet visited in our solar system, with surface temperatures reaching minus 390 degrees.

How the Planets Move


From our home planet, the Sun appears to circle Earth, rising in the east and setting in the west. But Earth really is the body that is moving. To understand the movements of the night sky, keep in mind Earth's two main motions: It circles the Sun, going around once in about $365\frac{1}{4}$ days (a year), and it spins on its axis, making one complete rotation in 24 hours (a day).

From night to night, each planet shifts its position slightly eastward in relation to the stars, traveling "through" various constellations as it circles the Sun. Mercury, Venus, and Mars show this motion most clearly. Distant Jupiter and Saturn travel so slowly, it's harder to see their eastward shift against the stars.

Earth's counterclockwise spin—from west to east—makes the sky overhead seem to rotate in the opposite direction, from east to west.

The retrograde motion of the planets puzzled early astronomers, who named these bodies “planets,” which in Greek means “wanderers,” because of the complicated ways the planets move.

Sometimes, a planet exhibits *retrograde motion*—it appears to stop its eastward drift and loop back toward the west before resuming its normal west-to-east movement. For example, because Earth takes less time to orbit the Sun than outer planets like Mars, Jupiter, and Saturn, it occasionally overtakes an outer planet, like a fast car passing a slower one on the highway. As Earth approaches, the other planet appears to stop its eastward drift and loop back toward the west. As Earth swings past the planet, we see the planet resume its normal west-to-east drift in the night sky.



As Earth overtakes a planet that is farther from the Sun, the planet's movement appears to change direction. This phenomenon is known as retrograde motion.

Observing the Planets

Never look at the Sun to locate Mercury, as blindness could result. Make sure the Sun's disk is completely below the horizon before looking in its direction.


Five planets are visible from Earth with the unaided eye: Mercury, Venus, Mars, Jupiter, and Saturn. Uranus may be dimly viewed on very clear nights, but only with powerful telescopes can we view Neptune even faintly.

The Inferior Planets

Mercury and Venus orbit the Sun inside Earth's orbit, so they are known as *inferior* planets. When these planets are between Earth and the Sun, the side where they are lighted faces away from us, making them difficult to see. Behind the Sun, they are out of sight. The best time to see these planets, therefore, is when they are on either side of the Sun.

Because Mercury is so close to the Sun, it is never seen in a fully dark, nighttime sky. It is visible only in bright twilight, either very low in the western sky just after sunset or very low in the east just before sunrise.

Late winter and early spring are usually the best times for observers in the northern hemisphere to spot tiny Mercury in the evening sky, about half an hour after sunset. To find it in the early morning sky, about half an hour before sunrise, look in the direction of sunrise during the late summer and fall.



Mercury's phases

Venus also appears only in the early evening or early morning sky, but it is much easier to see than Mercury. Sometimes called the morning or the evening star, diamond-white and brilliant Venus is the brightest planet as seen from the Earth. If Venus is in the evening sky, you can't miss it.

Because different parts of their sunlit areas are visible from Earth at different times, Mercury and Venus have phases similar to the Moon's. At different times, they appear full, *gibbous* (three-quarters full), half-full (like a first-quarter or last-quarter moon), as a crescent, or anywhere in between.


When Mercury or Venus is on the same side of the Sun as Earth, the planet's unlit side faces Earth. As the planet moves around the Sun away from Earth, its sunlit side becomes visible as a thin crescent and grows gradually as the planet moves along. When Mercury or Venus is near the other side of the Sun, we can see almost all of its sunlit face—the full phase.

A conjunction occurs when two or more planets appear near each other in the sky. Major conjunctions—when several planets align—are rare celestial events that may occur decades apart.

The Superior Planets

The planets beyond Earth's orbit are known as the *superior* planets. Unlike the inferior planets, which are always close to the Sun in our sky, the superior planets can appear anywhere in the sky on the ecliptic, and they always appear full. Mars, Jupiter, and Saturn can be seen high in the night sky, long after the sun has set.

The best time to view a superior planet is when it is in *opposition*—when Earth is directly between it and the Sun. When a planet is in opposition, it rises at sunset and sets when the Sun rises.



The zodiac is centered on the *ecliptic*, the path the Sun appears to follow among the stars.

Tracking Visible Planets

All the planets travel within the narrow band of the zodiac, although they wander around a bit, so they can be easy to miss. Even if you have a pretty good idea about the location of the zodiac, you may still have trouble finding the planets. For help, look at a sky map, which you can download from the internet (with your parent's permission) or find at the local library. Here are a few tips on recognizing the planets.

Mercury. This small planet is hard to spot, but it is visible without a telescope, either just after sunset or just before dawn, always in the general direction of the Sun and never far from it.

Venus. After the Sun and Moon, Venus is the brightest object in the sky. It can sometimes be seen in daylight, but it is never more than three hours ahead of or behind the Sun.

Mars. Mars is called the Red Planet with good reason, but its brightness varies. Check its position in an astronomical almanac or guidebook.

Planets do not twinkle like stars because they have a measurable disk, and more light reaches Earth. So, the steadiness of the light is a good indication that you are looking at a planet.

Jupiter. Among the planets, Jupiter is second only to Venus for brightness. Its progress across the sky can seem so slow that you might mistake it for a star unless you track its movement for several nights.

Saturn. You will not see Saturn's famous rings with the unaided eye, but the planet is bright enough to find. Like Jupiter, Saturn takes a long time to show movement, so consult an almanac or guidebook before scanning the heavens for it.

Uranus. This distant planet is barely visible to the naked eye under even good viewing conditions. It moves very slowly, so to the unaided eye it looks like an average star. Through a telescope it appears pale green.

Pluto was discovered in 1930 by U.S. astronomer Clyde Tombaugh. In 2006, the International Astronomical Union decided Pluto should no longer be considered the ninth planet in our solar system because it meets only two of the three criteria for an object to be considered a full-fledged planet. Although Pluto orbits our Sun just like the other planets, and it has enough gravity to pull itself into a spherical (round) shape, it is now referred to as a "dwarf" planet because it is smaller and cannot clear other objects out of its path. Remarkably, even with its small size, we have learned through the use of the Hubble Space Telescope that Pluto has five moons of its own.

Asteroids, Comets, and Meteoroids

Other members of the Sun's family include asteroids, comets, and meteoroids—sometimes considered “cosmic debris.”

Asteroids



Ida the asteroid

Asteroids are minor planets of the Sun. Estimates of their number range into the millions. Some are only a few feet in diameter. The largest, Ceres, is about 580 miles across.

Most asteroids orbit the sun in the huge expanse of space between Mars and Jupiter known as the asteroid belt or main belt. Sometimes the gravity of massive Jupiter changes the orbit of an asteroid so that it crosses the orbits of the inner planets. Asteroid fragments strike Earth every day, and scientists believe several large asteroids have slammed into Earth over the years.

Comets

Some scientists believe that a huge asteroid many years ago struck near Mexico's Yucatán Peninsula, setting off a chain of events that doomed Earth's dinosaurs to extinction.

Comets are often described as dirty snowballs because they are largely made up of ice embedded with rock and dust particles. Most comets travel in long, elliptical orbits around the Sun.

When a comet nears the Sun, the intense heat causes the gases and particles that were frozen together to “melt” and vaporize. The vapor forms a “tail” that may be millions of miles long.



Comets that are bright enough to be seen without a telescope are so rare that only two or three comets may be visible from Earth every 10 or 15 years.

Meteoroids, Meteors, and Meteorites

These three are different aspects of the same members of the solar system. They start as *meteoroids*—stony or metallic materials in space. Some are microscopic, others are huge.

Every day, Earth's atmosphere collides with millions of meteoroids that burn in the tremendous heat created by the collision. If a meteoroid is large enough, the collision is visible as a streak of light across the sky—a *meteor*, or shooting star.

Some meteoroids survive the flaming fall to Earth without burning up completely. Those that make it to the ground are *meteorites*. Most meteorites are small, but the largest meteorite ever discovered weighs about 66 tons. It fell on a farm in South Africa.



The Willamette Meteorite was found in 1902 in Oregon's Willamette Valley. It weighs 15.5 tons and measures 9.8 feet long and is the largest meteorite ever found in the United States.

Meteor Showers

Some meteoroids are believed to be clusters of comet fragments that continue to orbit along the comet's old path, creating showers that blaze through the sky as they pass through Earth's atmosphere. Most meteor showers are strongest in the early morning hours, between midnight and 4 a.m. Some noteworthy meteor showers, named for the constellations from which they come, are listed here.

Shower	Date of Maximum
Quadrantid	January 3–4
Eta Aquarid	May 6–7
Perseid	August 12–13
Leonid	November 17–18
Geminid	December 12–13



The Moon—Our Nearest Neighbor

The Moon was humankind's first target in exploration beyond Earth. People first studied the Moon with just their eyes, then with Earth-based telescopes, then with robots in space. In 1968, explorers flew to the Moon and studied it from orbit. The following year, U.S. astronauts landed on the Moon and brought back rock and soil samples.

Six times between 1969 and 1972, U.S. astronauts explored the lunar surface and left behind scientific instruments to help scientists learn more about Earth's natural satellite. A few robotic probes have studied the Moon in the decades since, but the Moon remains a fascinating place with many unanswered questions.

The Face of the Moon

The Moon's surface is highly uneven, covered with mountains, plains, and valleys, and pitted with craters. The Moon is barren, without vegetation. It has no atmosphere and there are only traces of water. There are no clouds, no fog, and no snow to block our view of its surface.

These conditions also cause extreme surface temperatures. During a typical lunar day, which lasts two weeks, the Moon's surface becomes much hotter than water's boiling point on Earth. During lunar nights, which also last two weeks, the surface temperature plummets to more than 300 degrees below freezing.


The soil samples astronauts brought back from the Moon lack many essential minerals, but they are not poisonous to plants. Researchers mixed a tray of lunar soil with dirt and grew plants in it.

Even though one side of the Moon always faces away from Earth, both the near side and the far side of the Moon receive the same amounts of sunlight. There really is no "dark side of the Moon."

Mountains, Seas, and Craters

Go out at night and take a good look at the Moon. You will see lighter areas and darker areas. The darker areas were called *maria* (MAHR-ee-ah), or seas, because the astronomers who pointed the first crude telescopes at the Moon in the 17th century thought these were large bodies of water. The lighter areas were called *terrae* (TEH-ray), or lands. Look at the Moon through binoculars and you can easily see how early astronomers might have reached that conclusion.

Seas on the Moon




*This big plain is called an “ocean” because of its size—1,550 miles across.

We now know that the bright lunar lands are rough, mountainous highlands. The dark seas are broad, level plains. They are lowlands filled by ancient lava flows.

Millions of bowl-shaped cavities or pits, called craters, pockmark the Moon’s surface. Some craters span more than 100 miles wide. Some are thousands of feet deep. The Moon’s broadest craters often are called *walled plains* because their sides are much like steep cliffs rising up to mountain height.

Most lunar craters carry the names of astronomers, explorers, or ancient scientists and mathematicians, but several craters were named for mythological figures.



Craters and mountains on the Moon

Your sketch of the Moon for requirement 6a does not have to show a full moon, but you should select a phase that shows at least five seas and five craters. Draw the contours of visible features and label seas and prominent craters. If you have trouble identifying craters, most public libraries have astronomy books with lunar maps.


The Moon's Phases

Due to the rotation of the Earth, the Moon appears to travel westward since we see it rise in the east and set in the west. If you watch the Moon's position against the background stars, you will see that it also moves slowly eastward due to its orbit around the Earth. This eastward motion means that the Moon rises about 50 minutes later each day.


The Moon takes an average of $27\frac{1}{4}$ days to make one trip around Earth. But this is not the actual time from new moon to new moon. Earth also is moving, traveling in its orbit around the Sun. It takes the Moon about $2\frac{1}{4}$ more days to return to a position directly between Earth and the Sun, so the average time for the Moon to make one complete circuit from new moon to new moon is about $29\frac{1}{2}$ days.

The Sun always illuminates one half of the Moon, but, as the Moon orbits the Earth, we cannot always see all of the illuminated face. Just after new moon we see a slender crescent. At the first-quarter phase we see the Moon half-illuminated. During the following week, a gibbous (more than


A moon growing to full is said to be *waxing*. As it shrinks from full to new, it is *waning*.




NEW MOON




FULL MOON




WAXING
CRESCENT




WANING
GIBBOUS




FIRST
QUARTER



LAST
QUARTER



WAXING
GIBBOUS



WANING
CRESCENT

Even as it moves through its phases, the Moon is always a big rocky ball, roughly 2,200 miles across. Its appearance changes because half of the lunar surface always faces the Sun, but that lighted portion does not always face Earth.

half-illuminated) moon grows larger still. At full moon, the Moon is opposite the Sun in the sky so we see a completely illuminated disk. After the full moon, it becomes a gibbous moon, then shrinks to half-illuminated as a third-quarter or last-quarter moon, then becomes a crescent moon again. Finally it returns to the new-moon phase, where it is invisible in the Sun's glare.

The Moon's position in the sky is slightly different each night, and it rises later each night, too. With each day, the Moon rises an average of about 50 minutes later, and it drops about 13 degrees farther behind in relation to the Sun.


One of the easiest ways to sketch the phases and daily positions of the Moon is to start your observations when the Moon is somewhere in the southwestern sky. On your first night of observing, draw a couple of the constellations that are visible. Next, draw in the Moon and write the date beside it on your sketch. At the same hour on other nights, add the Moon's new position.

The Moon's Orbit

The Moon orbits Earth in an ellipse, or slightly flattened circle, at an average speed of 2,300 miles per hour. Its average distance from Earth is 238,857 miles.

If the Moon traveled in a straight line, it would leave orbit and shoot off into space. The force that holds the Moon in its orbit is *gravity*, the same force that pulls a ball to the ground when you drop it. Because the Moon is much nearer to Earth than any other body, its motion is mainly determined by Earth's gravity, and it continuously "falls" around Earth.

Gravitational attraction exists between every pair of objects. So just as Earth's gravity tugs on the Moon, the Moon's gravity pulls on Earth with an equal force. That pull causes Earth to waver slightly as it revolves around the Sun. Lunar gravity also strongly affects Earth's ocean tides.



The Moon's orbit

Remember that Earth's surface is curved. If you could throw a ball fast enough so that it would land thousands of miles away, Earth would curve away from the speeding ball just as fast as the ball was dropping. When that happened, the ball would keep on falling and never strike the ground—it would be orbiting our planet.

The Moon travels faster at its *perigee*, when it is closest to Earth—221,000 miles away. It travels slower at its *apogee*, when it is farthest from Earth—about 253,000 miles away.

The *orbital plane* is the imaginary flat surface (the plane) that Earth marks out in space as it orbits the Sun. With few exceptions, all the other planets orbit in or near this same plane.

NASA has a helpful website devoted to eclipses past, present, and future. Go to eclipse.gsfc.nasa.gov/eclipse.html.

The Moon's Role in Eclipses


An eclipse happens when the Moon moves directly into line with Earth and the Sun. Eclipses can occur only at new moon or full moon, and then only when the Moon is crossing Earth's orbital plane. Eclipses usually occur several times a year.

There are two types of eclipses: solar and lunar.

A solar eclipse can be a dramatic event. It happens when the Moon passes directly between Earth and the Sun, so that the Moon's shadow strikes Earth. During a total solar eclipse, the Sun is completely blotted from view for everyone in the path of the Moon's shadow. A partial solar eclipse is visible off to one side of the path of total eclipse. The Moon appears to take a bite out of the Sun instead of fully blocking it.



Solar eclipse




When the Moon is at a far point in its orbit, an annular eclipse may occur. In this case, the Moon appears smaller because it is farther away, and it blocks only the center of the Sun's disk, allowing sunlight to blaze all around its edges.

A lunar eclipse happens when the Moon moves directly behind Earth, passing into Earth's shadow. There are several types of lunar eclipses:

- **Partial lunar eclipse**—The Moon does not travel completely into Earth's shadow.
- **Penumbral eclipse**—The Moon is completely within the brighter, outer portion (penumbra) of Earth's shadow.

- **Umbral eclipse**—The Moon passes into the darkest inner portion (umbra) of the shadow.
- **Total lunar eclipse**—The Moon moves completely within the umbra. It is the rarest of lunar eclipses.


You don't need any special equipment to view a lunar eclipse. Lunar eclipses often go unnoticed, even though they are visible over a great area. Most people fail to notice a slight shadow, dimming, or discoloration of the full moon.



Lunar eclipse


Viewing a Solar Eclipse


Seeing a solar eclipse can be an unforgettable experience—but don't let that be the last thing you ever see. **Never** stare directly at the Sun, even for a few moments. Doing so can cause permanent blindness. *Projection is the safest way to observe solar activity.* However, when using the projection method, you should **never** look through the eyepiece.



If you don't have a telescope or binoculars, use the pinhole projection method. You will need two pieces of flimsy cardboard. Using a straight pin, make a small hole in the center of one piece. Hold this first piece of cardboard above the second one, lining the cards up with the Sun as shown. The light will project an image of the Sun's eclipse through the hole in the first card onto the second card.

One way to safely view the Sun's image is by projecting it through the objective lens (the large end) of a telescope or binoculars and holding a piece of paper under the eyepiece like a screen. An image of the Sun will display on the paper. **Never** look into the eyepiece.





The Sun and Stars

The Sun is a huge ball of hot gas about 93 million miles from Earth. It contains 99.86 percent of the mass of the entire solar system. If you imagine Earth as the size of a grape, then the Sun is about 5 feet high—the size of a small refrigerator.

The Sun is made mainly of hydrogen and helium. Energy is released at the Sun's core as hydrogen changes into helium during nuclear fusion reactions. Every second, the Sun converts 564 million tons of hydrogen into 560 million tons of helium. Four million tons of matter are converted into energy that the Sun then radiates into space.

The Sun produces enough energy every second to meet the needs of the whole world for more than 600,000 years.



The Sun From the Inside Out

The *photosphere* is the Sun's outer surface—the visible layer from which most of the Sun's radiation and light escape.

In the *convective zone*, energy is carried by rising and falling currents of hot gas called *convective cells*.

The *radiative zone* is the section where energy leaves the core.


The *core* is the center of the Sun, where nuclear reactions take place. The temperature at the core is close to 29 million degrees. The core occupies 2 percent of the Sun's volume but has 60 percent of its mass.

Above the photosphere lies the hot gas of the *chromosphere* ("sphere of color"), which is in violent motion. The chromosphere may be seen briefly during total solar eclipses as a reddish rim around the Sun.


Above the chromosphere is the *corona* ("crown"), which extends outward from the Sun in the form of the "solar wind."

Sunspots

Sunspots are cooler areas on the Sun (but still about 7,000 degrees F). They appear in the photosphere as pairs or groups of dark spots that can measure more than 20,000 miles across. When sunspots are numerous, the total energy coming from the Sun increases slightly. Hot material near a sunspot can burst out from the Sun in what is known as a *solar flare*.



Sunspots are not permanent features of the Sun's surface. They come and go in cycles, usually over a period of about 11 years.



Particles, radiation, and magnetic fields from solar flares bombard Earth.


Sunspots can cause static and drown out radio signals on Earth. If the energetic flow from the Sun is strong enough, it can cause power surges, damage sensitive electronics, and overload power systems on Earth.

The Sun and Earth's Weather


The Sun's energy drives weather and climate on Earth. It burns off Earth's moisture by evaporation, heats the atmosphere, and creates wind when one air mass becomes hotter than another. Clouds condense from water vapor that has evaporated from the oceans, and rain from these clouds returns water to the oceans.

The Reason for Our Seasons

Earth has seasons because of its tilted axis, which keeps the northern hemisphere tipped toward the Sun for half the year and away for the other half. The northern hemisphere gets more direct sunlight in summer, and less sunlight six months later in winter. Thus, our region receives and retains less heat from the Sun during the winter.



From our viewpoint on Earth, the Sun seems to rise in the east and set in the west. But that is actually true only on the first day of spring and the first day of fall. In the winter, the Sun rises southeast and sets southwest. During summer, it rises northeast and sets northwest.



The average distance from Earth to the Sun is 93 million miles.

Viewing the Sun

It bears repeating: **Never** view the Sun directly; your eyesight is too precious to risk. Telescope filters are not recommended because you are still looking in the Sun's direction, so part of your line of vision falls outside the telescope lens. Also, filters can shatter, leaving the eyes unprotected, which can blind the observer instantly. *View the Sun only by projection.*

Star Quality

Stars are divided into color categories, from hottest to relatively coolest: blue and blue-white, white, yellow, orange, and red. Stars are referred to this way because they appear to be these colors when observed through a telescope.

Blue and Blue-White Stars. The hottest of all stars appear blue or blue-white. Their temperatures range from 18,000 degrees F to more than 45,000 degrees F. An example of a hot, blue-white star is Rigel in Orion. Others include Altair in Aquila and Deneb in Cygnus (both are in summer constellations).

White Stars. White stars have temperatures from 13,500 degrees F to about 18,000 degrees F. Sirius in Canis Major—the brightest star in our sky (not counting our Sun)—is a white star. So are Polaris in Ursa Minor and Procyon in Canis Minor.

Yellow Stars. Yellow stars are cooler, at temperatures of about 9,000 to 13,500 degrees F. The Sun is a yellow star, and so is Capella in Auriga. For historical reasons, astronomers refer to these yellow stars using the letter “G,” and thus our Sun is a “G star.” Other yellow stars include Lambda in Sagittarius (top of the “teapot”) and Zeta in Cygnus (part of the swan’s wing).

Orange Stars. Orange stars are between 6,000 and 9,000 degrees F. Arcturus in Boötes is an orange star, as is Dubhe in Ursa Major.

Red Stars. Red stars, the “coolest,” are cooler than 6,000 degrees F. One of the brightest stars, Betelgeuse, is a red star. Antares in Scorpius also is a red star.

Brightness


The true brightness of a star is determined by its absolute magnitude or luminosity. A star’s brightness in our sky depends on both its true brightness and its distance from us. A star can be really bright and yet, if it is an enormous distance from Earth, it may appear faint in our sky.





The opposite also is true: A nearby star that is not terribly bright can shine brilliantly in our sky. Sirius, for example, is just nine light-years from Earth. It shines more brightly in our sky than does Rigel, which actually is a much brighter star but so far away (about 800 light-years) that it appears fainter to observers on Earth.

A bright star is not necessarily large, either. Some of the brightest stars are smaller than other, fainter stars.



Tools of the Trade

You can observe objects in the night sky with your unaided eye. However, with binoculars or a telescope, the images you see will appear brighter and larger.


Binoculars

Binoculars are perfect for stargazing. They collect more light than the human eye, so you can see many more stars with binoculars than with the unaided eye. They also improve the clarity and intensify the colors of the stars you see.

Unlike a telescope, binoculars allow you to use both eyes to view. Binoculars show an image the right way up, whereas telescopes show objects upside down. Binoculars are easy to transport and less expensive than many telescopes.

The increased light-gathering power of their wide lenses makes binoculars ideal for astronomy.

Because of their wide field of view, binoculars are perfect for studying the surface of the Moon, scanning the Milky Way, spotting Jupiter's large moons, and viewing star clusters. An ideal size is 10×50 , which means the image is magnified 10 times and the main lenses are 50 millimeters in diameter. The 8×40 and 7×35 are also good choices.



Standard binoculars




Some giant binoculars have lenses of 70 millimeters or more and magnifications of 15x to 20x. Wider lenses allow you to see more stars, but these giant binoculars are so heavy that they must be mounted on a tripod, like a telescope, for viewing.

Telescopes


Like binoculars, telescopes gather more light than the human eye. The two main types of optical telescopes—those that collect visible light—are *refracting* and *reflecting*. A third type, the *catadioptric*, combines features of the refractor and the reflector.

Principle of a refracting telescope



Refracting telescopes use a system of lenses. At the large end of the telescope, the objective or front lens collects and focuses light. The eyepiece, the smaller lens you look through, is at the other end. Refracting telescopes produce sharp images.

Refracting telescope



Principle of a reflecting telescope

In the reflecting telescope, a concave (bowl-shaped) mirror at the base of an open tube collects and reflects light to a second, smaller mirror near the top of the tube. The eyepiece magnifies the image that the mirrors have formed. The reflector is the most common type of telescope and a popular choice for backyard astronomers.



Reflecting telescope


Radio and X-Ray Telescopes

Radio telescopes pick up images that astronomers would not be able to see otherwise. Radio waves from space reflect off of the telescope's large bowl-shaped (dish) antenna, and the radio receiver picks up the signals. A computer converts the signals into images. Radio signals reveal details, including temperature and composition, of objects in space that give off radio waves.

X-ray telescopes help scientists determine the galaxy's hot spots. Cosmic X-rays—the invisible radiation emitted from very hot objects—captured by telescopes above Earth's atmosphere help scientists study dying stars, colliding galaxies, and quasars, extremely bright starlike objects that give off enormous amounts of energy.

NASA's most sensitive X-ray telescope, the Chandra X-ray Observatory, captures X-rays from the edge of the observable universe.

X-rays from distant galaxies are absorbed by Earth's atmosphere, making them impossible to study from the ground.



The dish antennas of radio telescopes are much larger than the lenses or mirrors of optical telescopes. The Green Bank Telescope in West Virginia is one of the world's largest radio telescopes. Its dish measures over 330 feet across.

Ultraviolet and Infrared Telescopes

Ultraviolet and infrared telescopes detect light outside the visible electromagnetic spectrum. A rainbow represents the visible spectrum, with infrared light beyond the red end and ultraviolet light beyond the violet end.

Ultraviolet telescopes help astronomers study the formation of stars and galaxies. Because most of the ultraviolet waves from the Sun are blocked from Earth's atmosphere, these telescopes have to be placed in space in order to capture the ultraviolet light from the Sun, stars, galaxies, and other objects in space.

Infrared telescopes detect infrared light emitted by objects in space. Scientists use this data to detect brown dwarfs and distant galaxies. Infrared telescopes in space can see the coldest and dustiest objects in the universe.

The Hubble Space Telescope is an example of an ultraviolet telescope. The solar-powered reflecting telescope in an observatory that circles outside Earth's atmosphere is the size of a school bus. Its sophisticated equipment is operated by remote control and can get spectacular images of space.



The Hubble Space Telescope captured this image of plumes of interstellar gas and dust, called "elephant trunks," in the Eagle Nebula, some 7,000 light-years from Earth.

Care of Binoculars and Telescopes

Following the manufacturer's recommendations and a few simple tips will help you keep your viewing equipment in good shape for many years.

- Handle binoculars and telescopes carefully and with respect. Repeated bumping, shaking, and dropping can cause the lenses to break or become misaligned.
- Keep your equipment clean. Keep lenses free from dirt, debris, fingerprints, and moisture. Use a cloth (such as a microfiber cloth) and cleaner made especially for lenses. Apply cleaner very sparingly to the cloth and never directly to the lenses.



- Keep dust from building up on lenses. Aside from normal cleaning, you may use a very soft brush (such as camel's hair), or compressed air if you take some precautions. You want to avoid any contact with the lenses and any liquid from the can. Before use, avoid shaking the can. Always hold the can upright and never tilted or directly overhead while spraying. Use a few short (a second or two) bursts of air; stop immediately if the can starts to feel cold.

- Keep water away. Protect equipment from the elements such as rain, snow, and—in particular—saltwater, which is corrosive. Repeated use in high humidity will also have a negative effect on equipment.
- In the field, use and wear the strap that comes with your binoculars to keep them safely around your neck.
- Store binoculars and telescopes in a cool (not cold), dry place, away from humidity and extreme temperatures, which will have a negative effect on the lens coatings and housing. Use the storage or carrying case and rubber eyecups (in the “up” position) that came with your equipment.

Whenever your binoculars or telescope are not in use, be sure to leave on the lens caps.



Other Instruments Used in Astronomy

A *spectrometer* separates light coming from an object into a pattern of colored lines, much like a rainbow, called a *spectrum*. Each element, such as hydrogen, oxygen, or carbon, gives off or absorbs particular colors of light. So the unique color pattern can be used to identify the elements that make up the object. The colors can be shifted slightly if the object is moving toward or away from the Earth. Thus, a spectrum can also be used to measure the speed with which objects are approaching or receding from us.

Astrophotography

Astronomical photography allows you to keep a record of your observations of the sky and to share your experiences with friends and family. Modern digital cameras are very sensitive to light, and objects appear brighter and clearer in photographs than through a telescope or binoculars. In addition, you can adjust and process digital images to bring out even more details and colors that you cannot see with the eye.

There are many ways to photograph the sky. You can capture images of the Moon, planets, stars, constellations, nebulae, and galaxies with a digital single-lens reflex (DSLR) camera, a film camera, a specialized astronomical digital camera, or even the digital camera in a smartphone.



Taking Images Through A Telescope

Taking images through a telescope will allow you to see more details than photographing the objects with a standard lens.

The Moon and planets are so bright that only a short exposure (usually much less than 1 second) is needed. It is possible to photograph the Moon and planets by holding the camera up to the eyepiece, but it can be difficult to get the object in the center of the camera and hold it still while you trigger the exposure.

A smartphone adapter or camera adapter will hold the camera still and keep the object centered. You could also mount the camera on a separate tripod and aim it into the eyepiece. A camera remote control (or the headphones on many smartphones) will allow you to trigger the exposure without shaking the camera.

You can connect a smartphone to a telescope using a special adapter that will keep the object centered and hold the camera still.


To photograph faint nebulae and galaxies, long exposures are needed. The telescope will need a drive to track the sky and keep the object centered. You can download astrophotography apps for smartphones that allow you to take long exposures or use the “B” shutter speed setting on a DSLR.

Taking Images of Stars and Constellations

It is also possible to use just a smartphone camera or digital camera without a telescope to take wide-field images of the stars and constellations. You can even capture the International Space Station passing overhead or an occasional meteor!

You can take a short exposure to see the brightest stars, or you can take a long exposure to see the stars trail across the sky. You can also set your camera to take a series of short exposures, one after another, and “stack” them to simulate a long exposure. A tripod is necessary to keep the camera steady during the exposure. If you don’t have a tripod, lean the camera against a steady object.

On a very dark night you can leave the shutter open for minutes—or even hours—to capture the stars’ movement or even a meteor.



Photos taken with a digital camera can be downloaded very quickly and easily to your computer.




As the stars move across the sky, longer exposures will give you star trails, a record of the path in the sky taken by the stars.

Processing the Image

After you have obtained the image, you can use photo-editing software to increase the brightness, change the contrast, and adjust the colors. You can use software to automatically align and stack many short exposures to simulate what you would have seen in a long exposure. Processing your image can bring out details that were not obvious at first look. But be careful. Too much processing can make an image look unnatural!

Astronomical Research Cameras

Today, almost all research in astronomy is done with digital detectors. Just like consumer-type digital cameras, they use a charge-coupled device (CCD) or a complimentary metal-oxide semiconductor (CMOS) device to record images. Professional research cameras are custom built. One of the largest, at the Vera C. Rubin Observatory in Chile, is the size of a small car and has 3.2 gigapixels (3.2 billion pixels)! In order to see objects 100 million times fainter than you can see with your eyes, the camera is cooled to a temperature of minus 150 degrees F.



The digital camera in the Vera C. Rubin Observatory in Chile has 189 individual CCD sensors and is more than 2 feet wide. Rubin Observatory is operated by the National Science Foundation's NOIRLab and SLAC (originally the Stanford Linear Accelerator Center).



Light pollution is a serious energy waste that concerns professional astronomers and amateur observers alike. In the United States, electricity that wastefully illuminates the night sky costs an estimated \$2 billion a year. The International Dark-Sky Association works to help provide energy-efficient options that direct light where it is wanted, not needlessly up into the sky.

Light Pollution

Light pollution is, very simply, too much artificial light concentrated in urban areas (areas around cities). The glare of streetlights, outdoor signs, parking-lot lights, dusk-to-dawn security lights, and outdoor fixtures around private homes creates glowing domes over cities and towns. It also can spoil your view of the night sky.

The Air Up There

Air currents also can affect observing. Turbulent air can ruin views of the Moon, planets, and faint nebulae. Windless nights are better for studying the Moon and planets. Other conditions that can affect observing are clouds, high or low humidity, and air pollution such as dust and chemicals. Faint stars may be visible only on clear, dark nights.

Seeing in a Bright Night

If light pollution interferes with your stargazing, try these tricks.

- Cover your head down to your shoulders with a dark, opaque (nontransparent) cloth. This will help prevent light from streetlights, passing cars, and the glowing sky itself from ruining your night vision.
- Observe the skies as late at night as you can. As businesses close and people go to bed, less light pollution will intrude on your stargazing.
- Look for the bright objects. Urban sky glow will not keep you from seeing Sirius, Vega, Spica, Aldebaran, or similarly strong shiners. You should also be able to spot the brightest star clusters—the Pleiades, Hyades, and Beehive clusters, for example—and the spectacular Orion Nebula.

During the last
few thousand feet
of its journey to
Earth, starlight that
has traveled
trillions of miles
through deep
space gets
washed out by
bright city lights.





You and Astronomy

Plan activities to have some fun with your newfound knowledge of astronomy. Invite friends, and share your knowledge with them.

Visit a Planetarium or Observatory

Check the activity schedule of a planetarium, science museum, or observatory in your area and round up some friends to visit. While there, pay close attention to the presentations and displays. At an observatory, find out about the telescopes in use. Most of all, ask lots of questions! The people on staff not only know about astronomy, they know about practical observing in your area.

Plan an Observing Session

Develop an observing plan for a session when you will set up a telescope or observe with binoculars. Assemble the charts and guidebooks you will need. Prepare a red-filtered flashlight so you can view your charts without impairing your night vision. Get a notebook or create an observing log.

Be sure to check the weather forecast and dress appropriately. Even in mild seasons like the spring, nights can get cold. It pays to have extra layers as well as gloves, a scarf, and a hat.



Scout out the area beforehand to make sure you are aware of any conditions or potential hazards on the ground that you might miss when you come back at night. Before leaving, let someone know where you will be and when you will return.

Throw a Star Party

Plan to do everything you did for your own observing session, plus more. Find a good date and a place with plenty of room for the number of people you want to invite. Considering the time of year and how far north you live, develop a list of space objects you wish to find during the star party.

Consider inviting Scouts in your troop, your neighbors, family and friends, classmates, and teachers. When you invite them, be sure to cover such subjects as weather and proper dress, observing etiquette, and safety. Encourage those who have binoculars to bring them.



Contact a local astronomy organization or club to volunteer to help with a public star party. If you are unsure whom to contact, ask at a college in your community.


Moon Observation

Date: _____ Time: _____


Telescope: _____ Power: _____

Crater(s): _____

Notes: _____



Write a summary of your observing session and share it with your counselor.



Careers in Astronomy

If you have enjoyed earning the Astronomy merit badge, you might want to consider a career in the science. To be an astronomer, you must be observant, logical, imaginative, intuitive, and curious—your goal is nothing less than to understand the nature of the universe. It helps to be patient and to have the determination to stick with a difficult problem or long project until you have seen it through, which can take years.

Astronomers typically are math and science whizzes, skilled with computers. You will need to take high school calculus or precalculus, physics, chemistry, and computer science to prepare for college astronomy.

Reading, writing, and speaking skills also are important. Astronomers give talks at professional meetings and write papers for scientific journals. So, when choosing your courses, do not neglect language, communication, and social studies.

Where Astronomers Are Found

Most astronomers teach at colleges and universities and also do some research in a particular area of astronomy, such as planetary science, solar astronomy, or the study of stars or galaxies. Many who teach astronomy may work in the physics department of their university. Often, those who teach astronomy also teach physics.

About a third of professional astronomers have careers in federal government or at government-supported national observatories and laboratories. About 10 percent of astronomers are employed in business or private industry like the aerospace field.

What Astronomers Do

Observational astronomers design and carry out observing programs with a telescope or spacecraft, but they do not spend all their time looking through the eyepiece of a telescope. They

typically spend only 10 to 30 nights per year working at an observatory or getting observations from spacecraft. And even then, they are not looking up at the heavens. They are working at computers, analyzing and interpreting the information collected by the telescope and other instruments.


People interested in astronomy also may find careers as science librarians or teachers, science writers, planetarium or science museum directors, observing technicians or assistants, telescope operators, and optical engineers. Related career opportunities might be found in computing, image processing, and instrument design and building.



Professional astronomers such as theoretical astrophysicists may never work directly with observing equipment. They often do their research using supercomputers and complex computer models.

Space for Amateurs

Serious amateur observers make important contributions to astronomy. Amateurs are the chief comet-finders. They map storms on Mars and Jupiter, track the movements of asteroids, and discover supernovas. Their thousands of telescopes, computers, and electronic cameras far outnumber the comparatively few large university and government observatories. You might prefer to remain a hobbyist, where you can do real science and you get to decide how to spend your stargazing nights.





Astronomy Resources

Scouting Literature

Deck of Stars; Night Sky pocket guide; *Chemistry, Digital Technology, Geology, Nuclear Science, Photography, Radio, Space Exploration*, and *Weather* merit badge pamphlets

With your parent's permission, visit the Boy Scouts of America's official retail website, www.scoutshop.org, for a complete listing of all merit badge pamphlets and other helpful Scouting materials and supplies.

Books

- Consolmagno, Guy, and Dan M. Davis. *Turn Left at Orion: Hundreds of Night Sky Objects to See in a Home Telescope—and How to Find Them*. Cambridge University Press, 2011.
- Davis, Kenneth C. *Don't Know Much About Space*. HarperCollins, 2001.
- . *Don't Know Much About the Solar System*. HarperCollins, 2004.
- Dickinson, Terence. *The Backyard Astronomer's Guide*, 3rd ed. Firefly Books, 2008.
- . *Hubble's Universe: Greatest Discoveries and Latest Images*, 2nd ed. Firefly Books, 2017.
- . *NightWatch: A Practical Guide to Viewing the Universe*. Firefly Books, 2006.
- Hall, Allan. *Getting Started: Long Exposure Astrophotography*. CreateSpace Independent Publishing Platform, 2013.
- Harrington, Philip, and Edward Pascuzzi. *Astronomy for All Ages*, 2nd ed. Globe Pequot Press, 2000.
- Lambert, David. *The Kingfisher Young People's Book of the Universe*. Kingfisher Books, 2001.
- Lang, Kenneth R. *The Life and Death of Stars*. Cambridge University Press, 2013.
- Legault, Thierry. *Astrophotography*. Rocky Nook, 2014.
- Mellinger, Axel, and Ronald Stoyan. *The Cambridge Photographic Star Atlas*. Cambridge University Press, 2011.
- Moché, Dinah L. *Astronomy: A Self-Teaching Guide*, 8th ed. Wiley, 2014.
- Price, Fred W. *The Planet Observer's Handbook*, 2nd ed. Cambridge University Press, 2000.

Schaaf, Fred. *40 Nights to Knowing the Sky: A Night-by-Night Skywatching Primer*. Henry Holt and Co., 1998.

CDs, DVDs, and Videos

Amazing Universe III. Hopkins Technology, 1995; CD-ROM.

High Resolution Astrophotography. DamianPeach.com, 2015; DVD.

IMAX Cosmic Voyage. Warner Home Video, 1996; DVD.

NOVA: Secrets of the Sun. PBS Home Video, 2012; DVD and Blu-ray.

Wonders of the Solar System. BBC Home Entertainment, 2010; DVD and Blu-ray.

Magazines

Astronomy
21027 Crossroads Circle
P.O. Box 1612
Waukesha, WI 53187-1612
Toll-free telephone: 800-446-5489
www.astronomy.com

Sky & Telescope
P.O. Box 420235
Palm Coast, FL 32142-0235
Toll-free telephone: 800-253-0245
www.skyandtelescope.com

Organizations and Websites

The Astronomical League

9201 Ward Parkway, Suite 100
Kansas City, MO 64114
Telephone: 816-333-7759
www.astroleague.org

International Dark-Sky Association

5049 E. Broadway Blvd., #105
Tucson, AZ 85711
Telephone: 520-293-3198
www.darksky.org

National Aeronautics and Space Administration

300 E. Street SW, Suite 5R30
Washington, DC 20546
Telephone: 202-358-0001
www.nasa.gov

NASA Solar System Exploration: Eclipses

solarsystem.nasa.gov/eclipses

NASA Solar System Exploration: Planets

solarsystem.nasa.gov/planets

National Radio Astronomy Observatory

520 Edgemont Road
Charlottesville, VA 22903
Telephone: 434-296-0211
public.nrao.edu

National Science Foundation NOIRLab

(formerly the National Optical-IR Astronomy Research Laboratory)
950 North Cherry Ave.
Tucson, AZ 85719
Telephone: 520-318-8000
NOIRLab.edu

The Planetary Society

60 S. Los Robles Ave.
Pasadena, CA 91101
Telephone: 626-793-5100
www.planetary.org

Sky Maps

www.skymaps.com

Space Telescope Science Institute

3700 San Martin Drive
Baltimore, MD 21218
Telephone: 410-338-4700
www.stsci.edu

spaceweather.com

www.spaceweather.com

Acknowledgments

The Boy Scouts of America thanks William Cress, Beaver Falls, Pennsylvania, who is a longtime Astronomy merit badge counselor and an experienced amateur astronomer. We appreciate immensely his knowledge, advice, and support in helping to update the *Astronomy* merit badge pamphlet. He gathered an impressive group of fellow astronomers to assist in this effort, and we thank them here, as well:

- Jane Konrad, Pittsburgh Regional Center for Science Teachers and the Pennsylvania NASA Educator Resource Center
- Dan Malerbo, Buhl Digital Dome, Carnegie Science Center, Pittsburgh, Pennsylvania
- Tim Manka, retired science instructor, an Astronomy merit badge counselor since 1966
- Jean Philpott, Buhl Digital Dome, Carnegie Science Center, Pittsburgh, Pennsylvania
- Pat Plunkett, Wheeling (West Virginia) Jesuit University
- Tom Reiland, founder, Wagman Observatory, western Pennsylvania

The Boy Scouts of America thanks Dr. Edward Murphy, professor of astronomy at the University of Virginia in Charlottesville for reviewing the *Astronomy* merit badge pamphlet and providing updates since 2016.

We are grateful to Marcus Valdez, Fanwood, New Jersey, who serves as assistant Scoutmaster of Troop 33, and to the Amateur Astronomers of William Miller Sperry Observatory at Union County College in Cranford, New Jersey, for their assistance with photography.

The Boy Scouts of America is grateful to Al Dorn and the Omaha Astronomical Society for their expertise, assistance, and guidance in updating requirements for the Astronomy merit badge and support in developing the 2004 edition. The Society's website, found at www.omahaastro.com, provides a wealth of information for the beginner amateur astronomer as well as those who have developed a more serious interest.

The Boy Scouts of America is grateful to the men and women serving on the National Merit Badge Subcommittee for the improvements made in updating this pamphlet.

Photo and Illustration Credits

Scott Berger, courtesy—pages 25, 54, 68, and 70

Digital Vision™ Astronomy & Space CD—page 51

Gemini Observatory/NOAO*/
AURA†/National Science
Foundation, courtesy—page 64

Goodshoot—page 5

NASA, courtesy—pages 27 (*Mercury*),
28 (*Earth*), 47, 59 (*telescope*), 74
(*two people*), and 76

NASA, ESA, and the Hubble Heritage
Team, courtesy—pages 29 (*Saturn*)
and 59 (*plumes*)

NASA, Jet Propulsion Laboratory,
courtesy—pages 27 (*Venus*),
30 (*Neptune*), and 36 (*asteroid*)

NASA, Marshall Space Flight Center,
courtesy—page 72

National Optical Astronomy
Observatory/Association of
Universities for Research in
Astronomy/National Science
Foundation, courtesy—
page 74 (*woman*)

National Radio Astronomy Observatory
(NRAO)/Associated Universities, Inc.
(AUI)/National Science Foundation
(NSF), courtesy—page 58

Orion Telescopes & Binoculars,
courtesy. Used by permission
—page 62

Jacqueline Orrell/SLAC National
Accelerator Laboratory/NSF/DOE/
Rubin Observatory/AURA—page 65

Royal Swedish Academy of Sciences,
courtesy—page 49 (*sunspots*)

SOHO Project, NASA Goddard Space
Flight Center, courtesy—
page 49 (*solar flares*)

Wikipedia.org/Dante Alighieri,
courtesy—page 37

All other photos are the property of
or are protected by the Boy Scouts
of America or are royalty-free stock
images.

Dan Bryant—page 6

John McDearmon—all illustrations on
pages 7, 12, 15, 17, 19, 21, 31–32,
34, 40–42, 44–45, 48, 50, and 55–57

Mary Oakes—pages 53 and 68

*National Optical Astronomy Observatory

†Association of Universities for Research in Astronomy